

FRUITS

APPLICATION: tree times during the season

first time during the budding stage 30 g/ha "profi" or 300 g/ha "green" 30 g/ha "profi" or 300 g/ha "green" second time prior to blossoming 30 g/ha "profi" or 300 g/ha "green" third time during picking

(to improve preservation in the storage)

Formulation		Dosing		Remark
identifier	mark	w h e n	how much	h o w
powder	WP	during the budding stage - first time -	two / tree times per 30g "profi" or 300g "green" dissolved in water per one hectare of plantation	dissolve 300g "green" into 4 to 5 liters, i.e. 30g "profi" into 1.5 to 2 liters of warm water which is to be divided to number of sprinklers necessary for spraying one hectare of plantation
		prior to blossoming - second time -		
		prior to or during picking - third time - (to improve preservation in the storage)		

Recommendation:

if there is an economic interest, spray prior to the appearance of color,


in other terms to spray simultaneously with application of other chemical protection preparations with 150 g/ha "green" (or 15g/ha "profi") for more successful recovery.



COMPLETELY HARMLESS FOR PEOPLE, ANIMALS (INCLUDING BEES) AND ENVIRONMENT; IT DOES NOT REQUIRE SPECIAL MEASURES OF HYGIENIC AND TECHNICAL PROTECTION *)

Ministry of Agriculture, Forestry and Water Management of the Republic of Serbia Decision № 321-01-02214/2019-11 dated February 10, 2020

*) Decision № 3/2–08–9291/02 dated 13/01/03 of Ministry of Labor, healthcare and social welfare of Federal Republic of Yugoslavia



APPLICATION

into plantation

- improves resistance to pests and diseases
- glossy leaves, deep green color (more chlorophyll)
- lusher blossoming with more successful fruiting
- better stand to climate extremes: morning frost / moisture / drought
- higher prevalence of class I fruit

effect on fruit

- increased varietal fruit color, by 2-3 nuances
- increased sugar and dry matter content in fruit
- favorable impact on the relation between the sugar content and total acids
- better preservation during transport and reduced rot in the storage

HIGHER FRUITAGE BY 5 TO 14% - RAISE THE CATEGORY OF THE FRUIT QUALITY (more proper for processing and storage!)

Application: as a rule 300 g/ha "green" (or 30 g/ha "profi"), three times during the season

First time: during the budding stage

Second time: spray prior to blossoming, dividing the quantity of powder dissolved in the already

prepared solution per hectare orchard of other preparations (not require a separate pass), respectively, the amount of water needed to spray one hectare.

spraying 5–10 days prior to picking (improve preservation in the storage) Third time:

Recommendation:

❖ if there is an economic interest, spray prior to the appearance of color, and with half the recommended dose every time you treat an orchard with any kind of pesticides

COMPLETELY HARMLESS FOR PEOPLE, ANIMALS (INCLUDING BEES) AND ENVIRONMENT; IT DOES NOT REQUIRE SPECIAL MEASURES OF HYGIENIC AND TECHNICAL PROTECTION *)

Dr Danica Gajic invents Agrostemin. As a unique preparation (it is neither a fertilizer, it is neither a pesticide, nor a phytohormone) with wide range of effects, based on researches, testings and verifications done by Scientific institutions and eminent scientists and based on proven effects in application, AGROSTEMIN® over 30 years of application in the agriculture in Yugoslavia has been classified as a standard agrotechnical measure, i.e. a preparation for improving the quality and increasing quantity of yield**).

AGROSTEMIN® is produced as per the Yugoslav Patent office document no. 32749. Considering the fact that there are also forgeries present on the market, with quality control we obtained the right to put the signature (& Danies Gy're) of the author-creator of Agrostemin - Dr Danica Gajic - on our packagings and, in such a way, to confirm their originality.

Decision № 3/2-08-9291/02 dated 13/01/2003 of Ministry of Labor, healthcare and social welfare of FR of Yugoslavia

Decision № 321–01–02214/2019–11 dated 10/02/2020 of Ministry of Agriculture, Forestry and Water Management of Serbia